

Association Of Candida Albicans Carriage And Dental Caries In Children

C Nandini¹, Nileshwariba Jadeja², Anil Patel³, Stuti Bajaj⁴, Shikha Jain⁵, Neetu Shah⁶

¹PhD scholar, Department of Oral Pathology and Microbiology, Gujarat University, Ahmedabad, Gujarat, India. Email: nandini.chaduvula@gmail.com

²Reader, Department of Oral Pathology and Microbiology, Karnavati school of dentistry, Karnavati university, Gandhinagar, Gujarat, India. Email: chavadanileshwari@gmail.com

³Reader, Department of Oral Pathology and Microbiology, Karnavati school of dentistry, Karnavati university, Gandhinagar, Gujarat, India. Email: anillpatel009@gmail.com

⁴Senior lecturer, Department of Oral Pathology and Microbiology, Karnavati school of dentistry, Karnavati university, Gandhinagar, Gujarat, India. Email: Stutibajaj@karnavatiuniversity.edu.in

⁵Reader, Department of Oral Pathology and Microbiology, AMC dental college and hospital, Ahmedabad, Gujarat, India. Email: shikha0207@gmail.com

⁶Lecturer, Department of Preventive and Community Dentistry, Karnavati School of Dentistry, Karnavati University, Gandhinagar, Gujarat, India. Email: neeturajeevshah@gmail.com

Corresponding Author:

Dr .C Nandini

Address: Karnavati University 907/A, Uvarsad, Gandinagar, Gujarat- 382422 India.

Email: nandini.chaduvula@gmail.com

DOI: 10.47750/pnr.2022.13.504.300

Abstract

Background: In the recent years *C. albicans* has been identified by researchers for having an important role in cariogenicity along with bacteria like *S. mutans* and lactobacilli, attributing to its acidogenic, aciduric and hetero-fermentive properties.

Objective: The present study evaluated the frequency, intensity, and species of oral Candidal carriage in children with primary and mixed dentition and to identify the association between oral Candidal carriage in children and dental caries.

Participants: 100 healthy children of age groups: group A: 4-7 years; and group B: 8-12 years, both caries active (divided into dmf/dmf+DMFT index 1-5, 6-10 and >10) and caries free were randomly selected for the study.

Study design: After a detailed intraoral examination, swab samples were taken from tongue and carious lesion for mycological examinations isolate *Candida* from their oral cavities in order to study the Candidal carriage in children with and without dental caries.

Results: Candidal carriage among children with high caries index was significantly higher than caries free children. 92.86% of the children from group A (4-7 years) having dmf index > 10 and 76.93% from group B (8-12 years) having dmf + DMFT index > 10 showed positive candidal carriage. Also, carriage of non albicans species of *Candida* increased with age.

Conclusion: The present study suggests a strong relation between Candidal carriage and dental caries in children. Further work needs to be done on larger samples recruited in community-based settings to substantiate the association of oral Candidal carriage with dental caries.

Keywords: Dental caries, *C. albicans*, Candidal carriage, cariogenicity.

INTRODUCTION:

Oral cavity is inhabited by over 400 species of microbes [1]. The percentage of Candidal colonization in oral cavity is found in good amount in immunocompetent persons and the most dominant is the *Candida albicans*. [2]. *Candida* spp. is endowed with “dimorphism”, in fact, a polymorphism [3-5] There has been an increasing interest in investigating the potential cariogenic properties of the oral fungal flora especially *Candida* and both presence and abundance of *Candida albicans* have been linked to high DMF indices [6,7].

In addition to streptococcus mutans and lactobacilli, *Candida albicans* has been frequently isolated from carious lesions in children. [8, 9]. The production of acid by acid producing organisms as the central etiological factor in the initiation of carious lesions has been considered in the ‘Miller’s acidogenic theory’. Lactobacilli along with other microbes like yeasts were mainly identified as possible candidates [10] and studies of oral *Candida* species prove their acidogenic and hetero-fermentative properties, especially in the presence of carbohydrates [11]. In a recent, ultrastructural study Dige and Nyvad in 2019 [12] showed co existence of cariogenic bacteria and fungi. Nikawa et al in 2003 [13] reported that *C. albicans* dissolved hydroxyapatite in a liquid culture at a 20-fold higher rate than *Streptococcus mutans*, despite a lower number of yeast cells. Klinke et al in 2009 [14] *Candida* remains active at a more acidic pH where streptococcus becomes inactive.

Children with high caries indices have shown *Candida* species in their saliva and it has been reported that the level of oral *Candida* species could be a risk factor in caries [15,16]. Hence, the present study was carried out to examine the occurrence of oral *Candida* in children with primary and mixed dentition and to identify the association between oral Candidal carriage in children and dental caries.

The present study was carried out in 100 school children, in age groups 4-6 years and 7-12 years, by performing mycological examinations to isolate *Candida* from their oral cavities with the aim to study the Candidal carriage in children with and without dental caries.

MATERIALS AND METHODS

PATIENT SELECTION:

In the present study, the correlation between dental caries and *Candida albicans* carriage in school children was determined by selecting 100 healthy children of age groups of : group A: 4-7 years; and group B: 8-12 years. Children who were having no relevant health histories were selected.

After the child was selected, his/ her demographic details were recorded and a thorough intraoral examination was done, which involved exclusion of any oral mucosal lesions, determination of the dmf or the dmf + DMFT index.

SAMPLE COLLECTION AND MYCOLOGICAL EXAMINATION

For mycological examination, a sterile mucosal swab was taken aseptically from dorsum of tongue and transported to the laboratory within 2 hours. The swab was inserted into 0.5 ml of sterile saline in a centrifuge tube, and a vortex mixer was used to rigorously mix for thirty seconds. 0.15 ml of the wash was spread onto three Sabouraud’s Dextrose medium containing chloramphenicol agar plates. The plates were incubated at 37 °C for 48— 72 h.

For collection of sample from the carious lesion, food debris and plaque was first removed thoroughly from the surface of carious lesion and carious dentine was excavated from the centre of lesion using an excavator. Care was taken to prevent contact of the excavator with the adjacent non carious enamel. The carious dentine was placed immediately into 0.5 ml of normal saline and was triturated using a sterile mortar and pestle. The triturated material was immediately inoculated on the Sabouraud’s Dextrose medium and incubated at 37 °C for 48-72 h and the formed colonies were identified as shown in the figure 1.

The colony forming units formed on each Sabouraud’s agar plate was counted using a semi automatic colony counter. Depending upon the number of colony forming units, they were divided into three groups: C.F.U. <10, 10-100 and > 100.

After that, germ tube test and chlamydospore formation on cornmeal agar were performed on the *Candida* isolates and an isolate with positive germ tube test and positive chlamydospore test was considered as *Candida Albicans* whereas an isolate with a negative germ tube and chlamydospore test were considered other candidal species. Sugar fermentation and assimilation tests were performed to detect the presence of species like *C. Tropicalis*, *C. Kefyr*, *C. Guillermondii*, *C. Parapsilosis*, *C. Krusei*, *C. Glabrata*. Sugar assimilation test result of *C. Albicans* is shown in figure 2.



Figure 1 - Figure showing candidal colonies formed on Sabouraud's Dextrose medium and incubated at 37°C for 48-72 h.




Figure 2 - Figure showing results of sugar assimilation test for C. Albicans species.

STATISTICAL ANALYSIS

The data was statistically analyzed using the SPSS statistical software. p- values, $p < 0.001$ was considered highly significant and $p < 0.05$ was considered as significant.

RESULTS:

In the present study, 100 children were selected and divided into two equal groups, 50 children each in Group A [4-6 years] Group B [7-12 years], out of which, 40 [80 %] children were caries active and 10 [20%] children were controls in each group as shown in the table 1.

As shown in the table 2, 54% of children in the group A and 42% children in group B showed Candidal carriage. In the group A, out of the 14 children with df-t index greater than 10, only 1 [7.14%] child was non-carrier and 13 [92.86%] children were carriers and from them, 11 [84.62%] children showed *Candida albicans*, 2 [15.38%] children showed *Candida tropicalis*. Similarly, among group B children, out of children with df-t +DMFT index greater than 10, 3[23.07%] children were non-carriers and 10 [76.93%] children were carriers and from them, 6[60%] children showed *Candida albicans*, 2 [20%] children showed *Candida tropicalis*, 1 [10%] child showed *Candida guillormondi* and 1 [10%] child showed *Candida parapsilosis*.

These findings suggested that the Candidal carriage was very strongly associated with dental caries activity showing a high statistical significance [$p < 0.001$]. Thus, suggesting an important role of *Candida* in causation of dental caries. Moreover, the age of the child may have an influence on the isolation of *Candida* species as the carriage of *Candida albicans* species decreased with age and there was a simultaneous increase in the carriage of non- albicans species in our study population.

Table 3 shows correlation between df-t +DMFT index and colony forming units [C.F.U.] isolated from tongue and carious lesion in group A and group B. The findings suggested that high caries indices are significantly associated with heavy Candidal colonization in both the carious lesions and tongue are strongly associated with causation of dental caries which are statistically significant [p< 0.05].

Table 1: Table 1: Table showing demographic details and caries activity status of the children in group A and group B.

GROUP	TOTAL NO.OF CHILDREN	SUBGROUP	NO. OF CHILDREN	SEX	
	NO. [%]		NO. [%]	MALES	FEMALES
				NO. [%]	NO. [%]
GROUP A [4-6 YRS]	50 [100]	CARIES ACTIVE	40 [80.00]	22 [55.00]	18 [45.00]
		CONTROLS	10 [20.00]	3 [30.00]	7 [70.00]
GROUP B [7-12 YRS]	50 [100]	CARIES ACTIVE	40 [80.00]	23 [57.50]	17 [42.50]
		CONTROLS	10 [20.00]	4 [40.00]	6 [60.00]
TOTAL			100 [100]	52 [52.00]	48 [48.00]

Table 2: Table 2: Table showing correlation between df-t + DMFT index and Candidal carriage in group A and group B.

GROUP	df-t +DMFT INDEX	TOTAL NO. OF CHILDREN	CANDIDA CARRIAGE		CANDIDA SPECIES			
			NON-CARRIERS	CARRIERS	C.ALBI CANS	C. TROPICALIS	C.GUILLO RMONDI	C.PARAP SILLOSIS
			NO. %	NO. %	NO. %	NO. %	NO. %	NO. %
GROUP A [50]	0	10 [20.00]	9 [90.00]	1 [10.00]	1 [100]	0 [0.00]	0 [0.00]	0 [0.00]
	1- 5	12 [24.00]	8 [66.67]	4 [33.33]	4 [100]	0 [0.00]	0 [0.00]	0 [0.00]
	6 - 10	14 [28.00]	5 [35.67]	9 [64.33]	5 +1* [66.67]	3 +1* [44.44]	0 [0.00]	0 [0.00]
	> 10	14 [28.00]	1 [7.14]	13 [92.86]	11 [84.62]	2 [15.38]	0 [0.00]	0 [0.00]
	TOTAL	50 [100]	23 [46.00]	27 [54.00]	22 [81.48]	6 [22.22]	0 [0.00]	0 [0.00]
GROUP B [50]	0	10 [20.00]	8 [80.00]	2 [20.00]	2 [100]	0 [0.00]	0 [0.00]	0 [0.00]
	1- 5	14 [28.00]	11 [78.60]	3 [21.40]	2 [66.67]	1 [33.33]	0 [0.00]	0 [0.00]
	6 - 10	13 [26.00]	7 [53.90]	6 [46.10]	6 [100]	0 [0.00]	0 [0.00]	0 [0.00]

	> 10	13 [26.00]	3 [23.07]	10 [76.93]	6 [60.00]	2 [20.00]	1 [10.00]	1 [10.00]
	TOTAL	50 [100]	29 [58.00]	21 [42.00]	16 [76.19]	3 [14.29]	1 [4.76]	1 [4.76]

* Shows one child with mixed carriage of *C. albicans* & *C. tropicalis*

Table 3: Table showing correlation between df-t +DMFT index and colony forming units [C.F.U.] isolated from tongue and carious lesion in group A and group B.

GROUP	DMFT INDEX	TOTAL NO. OF CHILDREN		C.F.U. ISOLATED FROM TONGUE				C.F.U. ISOLATED FROM CARIOUS LESION			
		NO.	%	0	< 10	10-100	>100	0	< 10	10-100	>100
				NO. %	NO. %	NO. %	NO. %	NO. %	NO. %	NO. %	NO. %
GROUP A [50]	0	10 [20.00]	9 [90.00]	0 [0.00]	1 [10.00]	0 [0.00]	10 [100]	0 [0.00]	0 [0.00]	0 [0.00]	
	1- 5	12 [24.00]	8 [66.67]	1 [8.33]	1 [8.33]	2 [16.67]	9 [75.00]	0 [0.00]	2 [16.67]	1 [8.33]	
	6- 10	14 [28.00]	5 [35.71]	0 [0.00]	8 [57.14]	1 [7.14]	8 [57.14]	0 [0.00]	3 [21.43]	3 [21.43]	
	> 10	14 [28.00]	1 [7.14]	2 [14.29]	6 [42.86]	5 [35.71]	5 [35.71]	0 [0.00]	6 [42.85]	3 [21.42]	
	TOTAL	50 [100]	23 [46.00]	3 [6.00]	17 [34.00]	7 [14.00]	32 [64.00]	0 [0.00]	11 [22.00]	7 [14.00]	
GROUP B [50]	0	10 [20.00]	8 [80.00]	1 [10.00]	1 [10.00]	0 [0.00]	10 [100]	0 [0.00]	0 [0.00]	0 [0.00]	
	1- 5	14 [28.00]	11 [78.57]	1 [7.14]	2 [14.29]	0 [0.00]	12 [85.71]	1 [7.14]	1 [7.14]	0 [0.00]	
	6- 10	13 [26.00]	7 [53.84]	0 [0.00]	6 [46.15]	0 [0.00]	9 [69.23]	0 [0.00]	2 [15.38]	2 [15.38]	
	> 10	13 [26.00]	3 [23.07]	0 [0.00]	7 [53.84]	3 [23.07]	4 [30.77]	0 [0.00]	5 [38.46]	4 [30.77]	
	TOTAL	50 [100]	29 [58.00]	3 [6.00]	15 [30.00]	3 [6.00]	35 [70.00]	1 [20.00]	8 [16.00]	6 [12.00]	

DISCUSSION:

Candida albicans is the most common fungus normally seen in human oral cavity. Several factors, such as persistence, dimorphism, acidogenicity, adherence and interference with host defense system work together to enhance the virulence of the fungi[13,17]. In the recent years there has been an increasing interest in the relationship between oral fungal flora, especially the *Candida* and dental caries. *Candida* is responsible for demineralization of tooth enamel and dissolution of hydroxyapatite, thereby causing caries[18, 19].

Therefore, study was performed to find the correlation between *Candida albicans* carriage and dental caries activity in children and to investigate the contribution of *Candida* in causation of caries. As it has been reported that a highest Candidal carriage is seen during the primary dentition stage [20], the study population was divided into two age groups : group A [4-6 years] consisted of children with primary dentition and the group B [7-12 years] consisted of children with mixed dentition.

The overall Candidal carriage rate of both the groups was of 48% which is quite similar to the findings of both Raja et al. [2010] and Moalic et al. [2001] who found an overall Candidal carriage rate of about 60%

[21,22]. In the present study, the Candidal carriage rate among the children in group A [primary dentition stage] was 54% and that in the children of group B [mixed dentition stage] was 42% which showed that Candidal carriage in the primary dentition stage was comparatively greater than mixed dentition stage but was statistically non significant and these findings were similar to the findings of RońCkiewicz et al. [2006] and Qi et al. [2005] [15, 20]. According to Kadir et al. [2005] and Starr et al. [2002] a lower Candidal carriage in elder children with mixed dentition and that may be due to the alterations in the oral hygiene habits and dietary habits may explain the reduction of Candidal carriage in this age group. [6, 23].

In the present study, 92.86% of children with high caries activity [>10] in group A and 76.93% children with high caries activity [>10] in age group B showed Candidal carriage with a significant difference between the Candidal carriage in the caries active children and caries free children in both groups which suggested a strong association between Candidal carriage and caries activity with a high statistical significance. These findings correlated with the findings of Signoretto et al. [2009] [24].

In the present study, *Candida albicans* was the principal *Candida* species isolated and represented 81.48% of all *Candida* isolated in group A and 76.19% in group B. The second species identified were *Candida tropicalis* [22.22% in group A and 14.29% in group B], with very low percentage of *Candida parapsilosis* and *Candida guilliermondii*. These findings were in accordance with those of Cortelliet al. [2006] [25].

In the present study, high DMFT indexes were strongly associated with high Candidal carriage. From the children in both group A and group B, with $df-t$ +DMFT indexes greater than 10, 78% and 64.29% children respectively showed Candidal carriage from tongue and 76.92% and 69.23% respectively showed Candidal carriage from carious lesion which were statistically significant. These findings were in accordance with those of Hossain et al. [2003] who suggested that the fungi residing and cultivating in the carious tooth may spread systemically to cause extraoral disease conditions.

The present study showed that *Candida* has a significant role in establishing a high caries activity. At present, the exact mechanism of caries production by *Candida* has not been elucidated, but it has been suggested by Klinke et al. [2009] that *Candida* facilitates the enamel and dentin demineralization by a complex process of acidification of the oral environment by different mechanisms. Firstly, by production of pyruvic acid, that has a low pKa of 2.36 and has greater acidification potential than lactic acid [pKa of 3.86] secreted by lactobacilli. The amount of pyruvic acid is greatest during the initial colonization periods, but gradually decreases later apparently because of its reutilization. Secondly, the cell membrane of yeast cells consists of H^+ -ATPase which expels protons out of the cells, causing acidification. Thirdly, an increased acidification occurs because of the carbon dioxide released during the metabolism in yeast bodies. [14] Also, abundance of *C. albicans* carriage creates an acidic environment in the oral cavity to promote dental caries. [27, 28]

CONCLUSION:

High loads of *Candida* may lead to occurrence of active dental decay in mixed dentition stage. Therefore, identification of high caries risk population by isolation of *Candida* is easier than isolation of other cariogenic microorganisms during epidemiological studies.

REFERENCES:

1. Savage DC. Mucosal microbiota. In: Ogra PL et al. eds. Mucosal immunology. New York: Academic Press, 1999.
2. Odds F C. *Candida* and candidosis. 2nd edn. Baillière Tindal, London. 1998.
3. Berman J., Sudbery P.E. *Candida albicans*: a molecular revolution build on lessons from budding yeast. Nat. Rev. Genet. 3, 918-930.
4. Nikawa H, Hamada T, Yamamoto T. Denture plaque –past and recent concerns. J Dent 1998; 26: 299–304.
5. Järvenpää A, Hietanen J, Rautemaa R, Sorsa T, Richardson M. *Candida* yeasts in chronic periodontitis tissues and subgingival microbial biofilms in vivo. Oral Dis 10: 106–112.
6. Kadir T, Uygun B, Akyuz S, Prevalence of *Candida* species in Turkish children: relationship between dietary intake and carriage. Archives of Oral Biology (2005) 50, 33–37.
7. Lozano Moraga CP, Rodríguez Martínez GA, Lefimil Puente CA, Morales Bozo IC, Urzua, Orellana BR. Prevalence of *Candida albicans* and carriage of *Candida non-albicans* in the saliva of preschool children, according to their caries status. Acta Odontol Scand. 2017;75:30–5
8. de Carvalho F.G., Silva D.S., Hebling J, Spolidorio L.C., Spolidorio D.M.P. Presence of mutans streptococci and *Candida* spp. in dental plaque/dentine of carious teeth and early childhood caries. Archives of Oral Biology (2006) 51, 1024–1028.
9. Marchant S, Brailsford SR, Twomey AC, Roberts GJ, Beighton D. The predominant microflora of nursing caries lesions. Caries Res 2001;35: 397–406.

10. Tanzer JM, Livingston J, Thompson AM. The microbiology of primary dental caries in humans. *J Dent Educ* 2001; 65: 1028–1037.
11. Thein ZM, Samaranayake YH, Samaranayake LP. Effect of oral bacteria on growth and survival of *Candida albicans* biofilms. *Arch Oral Biol* 2006; 51: 672–680.
12. Dige I, Nyvad B. *Candida* species in intact in vivo biofilm from carious lesions. *Arch Oral Biol*. 2019;101:142–146
13. Nikawa H, Yamashiro H, Makihira S, Nishimura M, Egusa H, Furukawa M. In vitro cariogenic potential of *Candida albicans*. *Mycoses* 2003; 46: 471–478.
14. Klinke T, Kneist S, de Soet J, Kuhlisch E, Mauersberger S, Forster A. Acid Production by Oral Strains of *Candida albicans* and *Lactobacilli*. *Caries Res* 2009; 43:83–91.
15. Rozkiewicz D, Daniluk T, Zaremba M.L., Cylwik-Rokicka D. Oral *Candida albicans* carriage in healthy preschool and school children. *Advances in medical sciences*. 2006; 51:187-190.
16. Coulter WA, Murray SD, Kinirons MJ, The use of a concentrated oral rinse culture technique to sample oral *Candida* and *Lactobacilli* in children, and the relationship between *Candida* and *actobacilli* levels and dental caries experience: a pilot study. *Int J Paediatr Dent* 1993; 3: 17–21.
17. Pereira D, Seneviratne CJ, Koga-Ito CY, et al. Is the oral fungal pathogen *Candida albicans* a cariogen? *Oral Dis*. 2018;24:518–526.
18. Maijala M, Rautemaa R, Järvelin A, Richardson M, Salo T, Tjäderhane L. *Candida albicans* does not invade carious human dentine. *Oral Diseases*. 2007; 13: 279–284.
19. Szabo B, Majoros L, Papp-Falusi E, et al. Studies on the possible aetiological role of different *Candida* species in pathogenesis of dentine caries by monitoring the calcium release from tooth particles. *Acta Microbiol Immunol Hung*. 2014;61:11–17.
20. Q. G. Qi, T. Hu, X. D. Zhou. Frequency, species and molecular characterization of oral *Candida* in hosts of different age in China. *J Oral Pathol Med* (2005) 34: 352–6.
21. Raja M, Hannan A, Ali K. Association of Oral Candidal Carriage with Dental Caries in Children. *Caries Res* 2010;44:272–276.
22. Moalic E, Gestalin A, Quinio D, Gest PE, Zerilli A, Le Flohic AM. The extent of oral fungal flora in 353 students and possible relationships with dental caries. *Caries Res* 2001; 35: 149-155.
23. Starr JR, White TC, Leroux BG, Luis HS, Bernardo M, Leitao J, Roberts MC. Persistence of *Candida albicans* carriage in healthy Portuguese schoolchildren. *Oral Microbiol Immunol* 2002; 17: 304–310.
24. Signoretto C, Burlacchini G, Faccioni F, Zanderigo M, Bozzola N, Canepari P. Support for the role of *Candida* spp. in extensive caries lesions of children. *New Microbiologica* 2009; 32: 101-107.
25. Cavalca Cortelli S, Campos Junqueira J, da Silva Faria I, Koga-Ito CY, Cortelli JR. Correlation between *Candida* spp. and DMFT index in a rural population. *Braz J Oral Sci* 2006; 5: 1007–1011.
26. Hossain H, Ansari F, Schulz-Weidner N, Wetzel WE, Chakroborthy T, Domann E. Clonality identity of *Candida albicans* in the oral cavity and the gastrointestinal tract of pre-schoolchildren. *Oral microbiology immunology* 2003; 18: 302-308.
27. Qian Du, Biao Ren, Jinzhi He, Xian Peng, Qiang Guo, Liwei Zheng, Jiyao Li, Huanqin Dai, Vivian Chen, Lixin Zhang, Xuedong Zhou, Xin Xu. *Candida albicans* promotes tooth decay by inducing oral microbial dysbiosis. *The ISME Journal* (2021) 15:894–908.
28. Fakhruddin KS, Samaranayake LP, Egusa H, Chi Ngo H, Panduwawala C, Venkatachalam T, Kumarappan A & Pesee S. *Candida* biome of severe early childhood caries (SECC) and its cariogenic virulence traits, *Journal of Oral Microbiology*, 12:1, 1724484, DOI: 10.1080/20002297.2020.1724484