

# MORPHOLOGICAL AND MOLECULAR PHYLOGENY OF ANOPELOCEPHALIDEAN CESTODE PARASITE FROM CAPRA HIRCUS(L) IN SOLAPUR DISTRICT (M.S.) INDIA

ASHA BHOGIL<sup>1</sup>, AMOL THOSAR<sup>2</sup>, SUNITA BORDE<sup>3</sup>

<sup>1,3</sup>Department of Zoology, Dr. Babasaheb Ambedkar Marathwada University, Aurangabad-431004.

<sup>2</sup>Department of Zoology, Vivekanand College, Aurangabad.

DOI: 10.47750/pnr.2022.13.S06.332

## Abstract

Cestode Parasite are collected from intestines of *Capra hircus* (L) in Solapur district (M.S.) India, were subjected to a complete parasitological examination. The present Cestode that is *Moniezia*Sp. differ from all known species is having the scolex quadrangular, mature segments nearly four times broader than long, number of testes, shape of ovary, number of Inter proglottidal gland and their arrangement. In morphological data the Cestode is identified as confirmed to be representing *Moniezia* species in mammalian host goat. Identity of Cestode Parasites was confirmed through amplification and sequencing of COI- 5P gene marker (by molecular characterization) Phylogenetic analysis of COI-5P gene marker of *Moniezia*Sp. showed that it is cladded with *Monieziaexpansa*.

**Keywords:** *Capra hircus* (L), COI-5P, Molecular identification, *Moniezia* Sp., Solapur.

## INTRODUCTION

Sheep and goat have a particular socio-economic importance, out of all the animals domesticated by human. In Asian countries, *Capra hircus*(L.) keeping is associated with poverty. It is predominately limited to rural, poor who have to keep *Capra hircus*(L.) for substance production and use milk in household their meat supplement the diet, hair and skin is a source of income *Capra hircus*(L.) are important livestock in India due to their great nutritional and economic value.

Parasitic infection cestode parasites belonging to genus *Moniezia* and *Stilesia*. The genus *Moniezia* was established by Blanchard, 1891. Recently Sunita borde et al., 2017 erected *Moniezia* (B) *bordae*, Jadhav V.M. et. al. 2018 erected *Moniezia*(B.) *madhavae* and lastly Amol Thosar, et al., 2020 erected *Moniezia*(B.) *shilae*. Recently Sunita Borde, et.al., 2021 erected two new species i.e., *Moniezia*(B.) *babai* and *Moniezia*(B.) *yamagutii*.

The gastrointestinal cestode parasites causes great loss in nutritional and economic value of host in different ways like lowering of fertility, reduction in food intake, reduction in milk and meat production, reduction in weight of host and may be mortality in heavily parasitized animals. Furthermore, parasites is appears to be a major factor for lowered productivity of Indian livestock. Hence the study of helminth (cestode) parasites needed today.

Aim of these study was to evaluate the phylogenetic position of the Cestode genera *Moniezia solapurensis*, Sp. Nov. of the Goat i.e. *Capra hircus*(L.) respectively from Solapur district (M.S.), India on the basis of morphological and molecular data (using molecular markers) among Anaplocephalidea.

The present communication deals with molecular and morphological identification of cestodes i.e. *Moniezia solapurensis*, Sp. Nov. from goats in Solapur district (M.S.)India.

## MATERIAL AND METHODS

Morphological identification: - Cestode parasites were collected from the intestine of goat in Solapur district (M.S.) India. These Cestodes preserved in 4% formalin and stained with Aceto-carmine or Harris hematoxylin, passed through various alcoholic grades, cleared in xylene, mounted in D.P.X. and drawings are made with the aid of camera Lucida. All measurements are given in millimeters, otherwise mentioned. The identification is made with the help of Systema Helminthum.

Molecular identification: -Cestodes intended for molecular analysis were fixed with 95% ethyl alcohol and transported to wet laboratory facility of Paul Hebert Centre for DNA Barcoding.

DNA isolated from tissue of cestode parasites using Promega wizard genomic Kit, quantified using Nano drop spectrophotometer (ND 1000, Thermo Corporation, USA) and quality checked on 1% Agarose gel electrophoresis. Gel was visualized using gel documentation system (Biorad Inc., USA). DNA diluted to have 100 ng/μl concentration and stored at -200C for further use. These DNA samples were processed further for molecular studies where we employed mitochondrial markers and microsatellite marker studies.

From mitochondrial genome we have selected two markers for amplifying Cytochrome oxidase- 1 gene. CO-I gene amplification was performed in 25 μl PCR reaction using Kappa bio systems PCR kit. The reaction constitute 12.5 μl of 10% trehalose, 5.5 μl H<sub>2</sub>O, 2.5 μl 10X PCR buffer, 0.4 μl MgCl<sub>2</sub>, 2.0 μl 2.5mM dNTPs, 0.5 μl 10mM primers (TGTAACGACGGCCAGTATTCAACCAATCATAAAGATATTGG(Hebert et al., 2004)) and 0.1 μl taq polymerase (5 units/μl) and 1 μl of 100 ng/ μl of template DNA.

The PCR thermal cycle conditions include an initial denaturation step of 2 min at 95°C, 35 cycles of 30 sec at 94°C, 40 sec at 52°C, and 1 min at 72°C, with final extension at 72°C for 10min. Amplified PCR products were visualized on 1.5% agarose gel as shown in figure No. 2

PCR products were processed for clean up to remove unincorporated nucleotide and residual primers using 0.25 μl Exonuclease-I (20 units/μl) and 0.5μl of Shrimp Alkaline Phosphatase enzyme (1unit/μl) followed by cycle sequencing reaction using BigDye® Terminator v.3.1 Cycle Sequencing Kit (Applied Biosystems, Inc.) with 16th fold dilution. During COI gene amplicon sequencing M13 primers were used whereas, for D-loop and Cyt b amplicon sequencing same PCR primers were used. The thermal cyclers conditions were an initial denaturation of 2 min at 96°C and 35 cycles of 30 sec at 96°C, 15sec at 55°C, and 4 min at 60°C. The Cycle sequencing is followed by sequencing clean-up by ethanol precipitation followed by dissolving template in HiDi formamide and bidirectional sequenced in ABI 3130 Genetic analyzer.

Sequences obtained for CO-I, region were aligned and assembled in Codon Code aligner v4.0.3 (Codon Code, Dedham, MA, USA) using Muscle algorithm and Mega 5 (Tamura et al., 2011) using Clustal- W algorithm.

Sequences have been deposited in NCBI Gen Bank under the following inclusive accession numbers for CO-I (JX024320 - JX423844).


Genetic divergence values for CO-I region were calculated using K2P parameters (Kimura, 1980). Publicly available cestode parasites sequences obtained from BOLD and GeneBank (NCBI) were also included in a separate analysis. Neighbor-joining trees (Saitou and Nei, 1987) of K2P distances were generated using Mega 5 (Tamura et al., 2011).

## RESULTS AND DISCUSSION

Morphological Description (Fig. 1)

Two specimens of the Cestode parasites were collected from the intestine of *Capra hircus* (L.) at Solapur district. (M.S.) India.

The Cestodes are long consisting scolex, neck and proglottids. Proglottids are immature, mature and gravid.



**Fig. 1. *Moniezia solapurensis*, Sp. Nov.**

The scolex is small in size, quadrangular in shape and measures, 1.36(1.170-1.37) in length and 2.28(2.47-2.08) in width. The Suckers are large, round in shape, four in numbers, arranged linearly and measures, 0.60(0.741-0.455) in length and 0.62(0.780-0.455) in width.

The neck is long wide anteriorly narrow and measures, 8.55(8.775-8.55) in length and 1.16(1.330-1.027) in width.

Mature proglottids are nearly four time broader than long, craspedote in shape each proglottids with a double set of reproductive organs and measures, 3.61(3.795-3.432) in length and 10.39(10.89-9.099) in width. The testes are small, oval to rounded in shape, 110-130 in numbers, distributed in longitudinal excretory canal of the segment and measures, 0.13 (0.165-0.099) in diameter. The vas-deference is thin, long tube, slightly curved measures, 2.57(2.838-2.31) in length and 0.10(0.132-0.066) in width. The Cirrus pouch is small, oval in shape, situated in middle margin of the segments and measures, 0.66 (0.759-0.561) in length and 0.49(0.825-0.165) in width. The Cirrus is thin, long, protruded curved inside the cirrus pouch and measures, 0.91 (0.990-0.825) in length and 0.10(0.132-0.066) in width.

The Ovary large in size, compact, round shape, two in numbers in each side of the segment and measures, 1.14 (1.419-0.858) in length and 1.34 (1.617-1.056) in width. The Vagina posterior to cirrus pouch, thin, long tube reaches to the Ootype and measures 2.13 (2.442-1.815) in length and 0.07(0.099-0.033) in width. The Ootype is small, rounded in shape, posterior to the ovary and measures, 0.31 (0.462-0.165) in diameter. The Genital pores are small, oval, marginal, middle in position and measures, 0.58(0.660-0.495) in length and 0.17(0.231-0.099) in width.

The Vitelline gland are large, rounded in shape, compact, post-ovarian and measures, 0.36(0.429-0.297) in length and 0.35(0.462-0.232) in width.

The Inter-proglottidal glands large, oval to rounded in shape, arranged in double row, present in between longitudinal excretory canals of the segment 20- 28 in numbers and measures, 0.26 (0.429-0.297) in diameter.

The longitudinal excretory canals are thin, present on both the sides of segments along the body lengths and measures, 0.26(0.396-0.132) in diameter.

The present worm *Moniezia solapurensis*, Sp. Nov., under discussion is having the scolex small, quadrangular, mature proglottids nearly four times broader than long, testes small in size, oval, 110-130 in numbers, cirrus pouch oval, ovary oval, vitelline gland post ovarian, inter proglottidal glands 22-28 in numbers.

- The present worm differs from *Moniezia (B) benedeni*, Moniez, 1879, Skrjabin and Schulz, 1937, which is having numerous proglottids broader than long, posterior proglottids fleshy, testes 500 in numbers, arranged in two groups, cirrus pouch short and wide, vas deferens with 2-3 coils, ovary compact, in the center of the segments, eggs well developed, inter proglottidal glands linear and close to the posterior margin of the segments, arranged transversely and reported from the Calves and Lambs.
- The present cestode differs from *Moniezia (B) pallida*, Monnig, 1926, which is having the uterus external, dorsal and ventrally over excretory canals, the inter-proglottidal glands varying in size and reported from the host horse in South Africa.
- The present parasite differs from *Moniezia (B) aurangabadensis*, Shinde, et al. 1985, which is having the scolex quadrangular, testes small, 1100-1200 in numbers, vas deferens coiled, cirrus pouch cylindrical, oval with some rounded acini, gravid proglottids broader than long, uterus reticulate, inter proglottidal glands 12-15 in numbers and reported from *Ovis bharal* (L.).
- The present tapeworm differs from *Moniezia (B) bharalae*, Shinde, et al, 1985, which is having testes rounded, 190-200 in numbers, vas deferens short, elongated, fusiform, genital pores bilateral, sub marginal, ovary compact, inter proglottidal glands arranged in two rows, small in size, 38-44 in numbers and reported from *Ovis bharal* (L.).
- The present form differs from *Moniezia (B) warananagarensis*, Patil, et al. 1997, which is having scolex large, globular, testes 300-320 in numbers, distributed throughout the proglottids, in single field, ovary indistinctly lobed with 13-15 short, blunt acini, transversely elongated, inter proglottidal glands, 56 in numbers, oval, medium in size, cirrus pouch medium, oval, transversely elongated, slightly obliquely placed and extend beyond longitudinal excretory canal.
- The present cestode differs from *Moniezia (B) kalawati*, Nanware, et al. 1999. Which is having squarish scolex, oval shaped cirrus pouch, testes small, oval, distributed throughout the segment, 172 in numbers, ovary medium, short, blunt acini, and 54 inter proglottidal glands in the inter segmental region, medium, oval either single or paired, irregularly arranged in the central width of the segments and leaving space on each lateral side.
- The present tapeworm differs from *Moniezia (B) murhari*, Kalse, et al, 1999, in having the scolex squarish, testes 405-415 in numbers, cirrus pouch elongated in the anterior region of the segments, ovary inverted horse shoe shaped, indistinctly bilobed each with numerous short, blunt, round, acini and inter proglottidal glands 63 in numbers.
- The present parasites differ from *Moniezia (B) caprai*, Pokale, et al, 2004, which is having the scolex is medium, squarish, with large four suckers, without rostellum, testes oval in shape, 255-260 in numbers, cirrus pouch is medium in size and ovary medium in size, kidney shaped.
- The present worm differs from *Moniezia (B) shindei*, Pawar, et al., 2004 in having scolex large, mature segments craspedote, testes 190-200 (195) in numbers, scattered all over segment and ovary a single mass, large, oval, cirrus pouch oval, elongated, in center of the segment and vitelline gland large, oval, internal to ovary.
- The present cestode differs from *Moniezia (B) hircusae*, Tat and Jadhav B. V., 2004 which is having scolex large, globular, mature segments big, craspedote, testes 168 in numbers, small, scattered in a single field, ovary large, oval, a single mass, in anterior half of the segment, inter proglottidal glands 14-15 in numbers, large, oval and cirrus pouch in anterior 1/3rd region of the segment.
- The present cestode differs from *Moniezia (B) rajalaensis*, Borde, et al. 2007, in having scolex large, globular, mature proglottids Squarish, Broader than long, testes 250-260 in numbers, medium, scattered throughout proglottids, ovary large, horse shoe shaped, inter proglottidal glands 31-32 in numbers, large, oval and cirrus pouch oval.
- The present cestode differs from *Moniezia (B) caprae*, Nanware S.S., 2010 in having scolex large, mature segment big, almost three and a half times broader than long, testes 84-85 in numbers, medium in size, oval in shape, ovary large, bilobed, inter proglottidal glands 40 in numbers, oval, rounded and cirrus pouch on each side.

- The present cestode differs from *Moniezia (B) govindae*, Padwal, et al., 2011 in having scolex large, globular, mature proglottids big, broader than long, testes 100-140 in numbers, medium, scattered throughout proglottids, ovary large, compact, nut shaped, inter proglottidal glands 40-42 in numbers, large, oval and cirrus pouch elongated.
- The present cestode differs from *Moniezia (B) babai*, Humbe, et al., 2011 in having scolex globular, mature segment four times broader than long, testes 190-220 in numbers, small in size, rounded in shape, ovary large, rounded, inter proglottidal glands 18-20 in numbers, oval, rounded and cirrus pouch on each side.
- The present cestode differs from *Moniezia (B) ovisae*, Humbe, et al., 2011 in having scolex broad anteriorly and narrow towards neck, mature segment two times broader than long, testes 155-165 in numbers, small in size, rounded in shape, ovary large, bilobed, inter proglottidal glands 32-35 in numbers, oval, rounded and cirrus pouch on each side.
- The present cestode differs from *Moniezia (B) osmanabadensis*, Humbe, et al., 2012 in having scolex globular, mature segment five times broader than long, craspedote, testes 170-200 in numbers, small in size, rounded in shape, ovary large, bilobed, inter proglottidal glands 38-40 in numbers, oval, rounded and cirrus pouch on each side.
- The present cestode differs from *Moniezia (B) devraoi*, Humbe, et al., 2013 in having scolex quadrangular, mature segment four times broader than long, testes 160-180 in numbers, small in size, rounded in shape, ovary large, bilobed, inter proglottidal glands 40-45 in numbers, oval, rounded and cirrus pouch on each side.
- The present cestode differs from *Moniezia (B) shegaonensis*, Barote, et al., 2013 in having scolex globular, mature segment four to five times broader than long, testes 190-220 in numbers, small in size, rounded in shape, ovary compact, inter proglottidal glands 20-25 in number, oval, rounded and cirrus pouch on each side.
- The present cestode differs from *Moniezia (B) shivajiraovae*, Barote, et al., 2014 in having scolex squarish, large in size, mature segment six to eight times broader than long, testes 84-95 in numbers, small in size, rounded in shape, ovary horse-shoe shaped, inter proglottidal glands 40-42 in numbers, oval, rounded and cirrus pouch on each side.
- The present cestode differs from *Moniezia (B) sureshi*, Ravi Solunke, 2015 in having scolex oval, quadrangular, mature segment four to five times broader than long, testes 180-185 in numbers, single field, unevenly distributed, ovary medium, horse-shoe shaped, in appearance having numerous short, blunt acini, inter proglottidal glands 18-19 in numbers, oval and cirrus pouch on each side.
- The present cestode differs from *Moniezia (B) jadhavii*, Amol Thosar, et. al., 2015 in having scolex squarish, mature segment craspedote, five times broader than long, testes 210-220 in numbers, small, oval to round, ovary horse-shoe shaped, compact, inter proglottidal glands 46-52 in numbers, arranged lineally in one or two rows, cirrus pouch small oval.
- The present cestode differs from *Moniezia (B) marathwadensis*, Shaikh Kalim, 2015 in having scolex quadrangular, mature segment five times broader than long, testes 125-130 in numbers, small, oval in shape, ovary compact with numerous blunt acini, inter proglottidal glands 50-52 in numbers, arranged lineally in one or two rows, cirrus pouch large, elongated, oval.
- The present cestode differs from *Moniezia (B) bhalchandrai*, Kalse A.T. et. al., 2016 in having scolex quadrangular, mature segment rectangular in shape, almost four and half times broader than long, testes 196-200 in numbers, oval in shape, ovary medium in size, inverted cup shaped, inter proglottidal glands 13-14 in numbers, oval in shape, highly muscular, single regularly and lineally arranged, cirrus pouch large, oval in shape.
- The present worm differs from *Moniezia (B.) bordeae*, Sunita Borde, et al., 2017 in having scolex quadrangular, mature segment nearly four to five times broader than long, testes 130-170 in numbers, spread in the medulla in between the longitudinal excretory canals, ovary bean shaped, small, forms concavity posteriorly, inter proglottidal glands 5-9 in numbers, arranged single row, cirrus pouch on each side and reported from *Ovis bharal* (L.).
- The present cestode differs from *Moniezia (B) madhavae*, Jadhav V.M. et. al., 2018 in having scolex quadrangular, mature segment near five times broader than long, testes 45-60 in numbers, medium in size, oval in shape, ovary distinctly bilobed, inter proglottidal glands 40-42 in numbers, oval in shape, cirrus pouch small in shape, curved.

- The present worm differs from *Moniezia (B.) shilae*, Amol Thosar, et al., 2020 in having the scolex quadrangular, mature proglottids nearly four times broader than long, craspedote in shape, testes small in size, oval to rounded, 180-210 in numbers, cirrus pouch oval, ovary large, compact, horse-shoe shaped, vitelline gland post ovarian, inter proglottidal glands 26-30 in numbers and reported from *Capra hircus (L.)*.
- The present worm differs from *Moniezia babai*, Sunita Borde, et. al., 2021 in having the scolex large, globular, mature proglottids nearly three times broader than long, testes small in size, round to oval, 150-170 in numbers, cirrus pouch oval, ovary butterfly shaped, vitelline gland post ovarian, inter proglottidal glands 20-25 in numbers.
- The present worm differs from *Moniezia yamagutii*, Sunita Borde et. al., 2021 is having the scolex large, globular, mature proglottids nearly three times broader than long, testes small in size, round to oval, 65-70 in numbers, cirrus pouch oval, ovary compact, rounded, vitelline gland post ovarian, inter proglottidal glands 28-33 in numbers.

The above differentiating characters are valid enough to erect a new species for these cestodes and hence the name *Moniezia (B.) solapurensis*, Sp. Nov. is proposed after the locality of host at Solapur district (M.S.) India

#### KEY TO THE SPECIES OF THE GENUS MONIEZIA, BLANCHARD, 1891

Mature segments broader than long	-	1
Mature segments Squarish	-	<i>M. (B.) pallida</i> , Monnig, 1926.
Mature segments Craspedote	-	2
1) Testes below 60 in numbers	-	<i>M. (B.) madhavae</i> , Jadhav V.M.et. al., 2018.
Testes between 65 -75 in number	-	<i>M. (B.) babai</i> , SunitaBorde, et.al., 2022
Testes between 80-100 in numbers	-	<i>M.(B.) caparae</i> , Nanware, et.al., 2010.
Testes in between 100-130 in number	-	3
Testes in between 130-150 in numbers	-	<i>M. (B.) kalawati</i> , Nanware, et al.,1999.
Testes in between 150- 175 in numbers	-	4
Testes in between 175-200 in numbers	-	5
Testes in between 200-300 in numbers	-	6
Testes in between 300 -400 in numbers	-	<i>M. (B.) warnanagarenisis</i> , Patil, et al., 1997.
Testes in between 400 -450 in numbers	-	<i>M. (B.) murhari</i> , Kalse, et al.1999.
Testes in between 450 -500 in numbers	-	<i>M.(B.) benedeni</i> , Moneiz, 1879,Skrjabin, et al., chulz, 1937.
Testes above 500 in numbers	-	<i>M. (B.) aurangabadensis</i> , Shinde, et al., 1985.
2) Inter proglottidal glands between10-15 in numbers -		<i>M. (B.)hircusae</i> , Tat and Jadhav, 2004.
Inter proglottidal glands between 16-20 in numbers	-	<i>M. (B.) babai</i> , Humbe, et al., 2011.
Inter proglottidal glands between 20-25 in numbers	-	<i>M. (B.) shegaonesis</i> , Barote, et al., 2013.
Inter proglottidal glands between 25-30 in numbers	-	<i>M.(B.) shilae</i> , Amol Thosar, et al., 2020
Inter proglottidal glands between 30-35 in numbers	-	<i>M. (B.) ovisae</i> , Humbe, et al., 2011.
Inter proglottidal glands between 35-40 in numbers	-	<i>M. (B.) osmabadensis</i> , Humbe, et al., 2012.
Inter proglottidal glands between 40-45 in numbers	-	<i>M. (B.) shivajiraovae</i> , Barote, et al., 2014.
Inter proglottidal glands above 50 in numbers	-	<i>M. (B.) shindei</i> , Pawar, et al. 2004.
3) Inter proglottidal glands between 20-30 in numbers -		<b><i>M.(B.) solapurensis,Sp.Nov.</i></b>
Inter proglottidal glands between 40-50 in number	-	<i>M. (B.) govindae</i> , Padwal ,et al. 2011.
Inter proglottidal glands above 50 in numbers	-	<i>M. (B) marathwadenesis</i> , Shaikh Kalim, 2015

- |   |   |   |
|---|---|---|
| 4) Scolex quadrangular  | - | <i>M.bordae</i> , Sunita Borde,et al.2017                           |
| Scolex quadrangular   | - | <i>M. (B.) devraoi</i> , Humbe, et al., 2013.                       |
| Scolex globular   | - | <i>Moniezia (B.) yamagutii</i> , Sunita Borde, <i>et.al.</i> , 2022 |
| 5) Inter proglottidal glands below 15                           | - | <i>M. (B.) bhalchandrai</i> , Kalse, et. al., 2016.                 |
| Inter proglottidal glands between 15-20 in numbers in two rows- |   | <i>M. (B.) sureshi</i> , Ravi Solunke, 2015.                        |
| Inter proglottidal glands between 20-45 in numbers-             |   | <i>M. (B.) bharalae</i> , Shinde, et al., 1985.                     |
| Inter proglottidal glands between 45-55 in numbers-             |   | <i>M.(B.) jadhavii</i> , Amol Thosar, <i>et al</i> ,2015.           |
| 6) Scolexsuarish  | - | <i>M. (B.) caparai</i> , Pokale, <i>et al.</i> , 2004.              |
| Scolex globular   | - | <i>M.(B.) rajalaensis</i> , Borde, <i>et.al.</i> , 2007.            |
| 7) Inter proglottidal glands between 40-50                      | - | <i>M. (B.) devraoi</i> , Humbe et.al, 2013                          |
| Inter proglottidal glands between 5-9 in numbers                | - | <i>M. (B.) bordeae</i> , Sunita Borde, et.al, 2017                  |

Molecular Data: -

A comparison of the partial sequences of the COI-5P gene of the present Cestodes with those of other Cestodes, in a Phylogenetic context, provided further support for placing this species as a new one within *Moniezia*solapurensis, Sp. Nov. thus confirming taxonomic conclusion based on morphological data.

In the phylogenetic trees (Fig.3) obtained by maximum parsimony analysis of the COI-5P sequence data set, Cestode species close to the species *Moniezia expansa* is clear with a maximum identity 99.84% (Table no.1). After partial COI-5P gene Sequence of *Moniezia* sample DNA sequences length is 628 bp (Fig.2)




Fig. 2. Amplification of COI-5P gene from sample




Fig. 3. Phylogenetic tree for *Monieziasolapurensis* Sp. Nov. using COI-5P gene sequence

Table No. 1:-Phylogenetic neighbors of *Monieziasolapurensis*, Sp.Nov. based on COI-5P gene sequence

Scientific Name	Max Score	Total Score	Query Cover	E value	Percentage Identity	Accession
<i>Monieziaexpansa</i>	1147	1147	99%	0	99.84%	OL689029.1
<i>Monieziaexpansa</i>	1147	1147	99%	0	99.84%	AB821382.1
<i>Monieziaexpansa</i>	1142	1142	99%	0	99.68%	AB821381.1
<i>Monieziaexpansa</i>	1136	1136	99%	0	99.52%	AB821383.1
<i>Monieziaexpansa</i>	1136	1136	99%	0	99.52%	AB821380.1
<i>Monieziaexpansa</i>	1136	1136	99%	0	99.52%	AB821379.1
<i>Monieziaexpansa</i>	1136	1136	99%	0	99.52%	AB821378.1
<i>Monieziaexpansa</i>	1136	1136	99%	0	99.52%	AB821376.1
<i>Monieziaexpansa</i>	1136	1136	99%	0	99.52%	AB821375.1
<i>Monieziaexpansa</i>	1136	1136	99%	0	99.52%	AB821374.1
<i>Monieziaexpansa</i>	1136	1136	99%	0	99.52%	LC102496.1
<i>Monieziaexpansa</i>	1131	1131	99%	0	99.36%	AB821377.1

## CONCLUSION

On the basis of morphological and molecular observations the above said parasite is erect as a new *Moniezia solapurensis* Sp. Nov. is proposed after the locality.

## ACKNOWLEDGMENTS

Author is thankful to the Head, Department of Zoology, Dr. Babasaheb Ambedkar Marathwada University, Aurangabad for providing the laboratory facilities during this research work and also thanks Paul Hebert Centre for DNA Barcoding and Biodiversity studies, Aurangabad laboratory for providing necessary facilities.

## REFERENCES

1. Amol Thosar, Asha Bidkar and Sunita Borde, (2020). A taxonomic study of a new cestode *Moniezia* (B) *shilae*, Sp. Nov. (Cestoda: Anoplocephalidae) in *Capra hircus* (L.) from Aurangabad district. Review of Research (Special issue). Volume-9, Issue-6. ISSN: 2249-839X. Impact Factor: 5.7631(UIF).
2. Amol Thosar, Ganesh Misal, Arun Gaware and Sunita Borde, (2015). Morphological and molecular studies of *Moniezia* Sp. (Cestoda: Anoplocephalidae) a parasite of the domestic goat *Capra hircus* (L.) in Aurangabad district (M.S.), India. Indian Journal of Applied Research. Vol.No. 5(8):10-14.
3. Amol Thosar, Rahul Khawal, Sushil Jawale and Sunita Borde, (2014). Some biochemical aspects of Anoplocephalidean Cestode Parasites in *Ovis bharal* (L.). The Ecoscan. Special issue, Vol. V:01-04, ISBN-0974-0376. IF- 1.86.
4. Barote, et al. (2013). On a new species of *Moniezia* Blanchard, 1891 (Cestoda: Anoplocephalidae) in *Ovis bharal* from Bhuldhana dist. (M.S.) India. Trends in Parasitology Research, volume 2(3):1-4 (Online).
5. Barote, et al. (2014). On a new species of *Moniezia* Blanchard, 1891 (Cestoda: Anoplocephalidae) in *Ovis bharal* from Bhuldhana dist. (M.S.) India. Trends in Parasitology Research, volume 3(1):1-4.
6. Blanchard, R. (1926). Sur les helminthes des primates antropoides, Mem Soc. Zool. France.:420-489.
7. Borde, S.N., Patil, P.S and Naphade, S.T. (2007). A new tape worm from the host *Capra hircus* at Rajala (M.S). Nat. J. Sci., 4 (3) :126-128.
8. Brand T Von, (1966). Biochemistry of parasites. Academic press, New York.
9. Chilton N.B. et. al. (2007). Genetic markers to distinguish *Moniezia expansa* from *Moniezia benedeni* (Cestoda: Anoplocephalidae) and evidence of the existence of cryptic species in Australia. Parasitol Res. 100:1187.
10. Felsenstein J. (1985). Confidence limits on phylogenies: An approach using the bootstrap. Evolution 39:783-791.
11. Hui Zhang et. al. (2008). Characterization of a *Moniezia expansa* ubiquitin-conjugating enzyme E2 cDNA. Mol. Biol Rep. 37:1585-1590.
12. Humbe Atul, S. N. Borde and Swati Jadhav, (2013). A New Mammalian Tapeworm *Moniezia devraoi* From *Capra hircus* at Amravati (M.S.) India. Weekly Science Research Journal.1(10):1-5.
13. Humbe, et al. (2011). On a new species of *Moniezia* *babai*, Blanchard, 1891 (Cestoda: Anoplocephalidae) from *Capra hircus* (L.) from Buldhana district (M.S.) India. International multidisciplinary Research Journal. 1(8): 01-03.
14. Humbe, et. al. (2011). Occurrence of a new mammalian tapeworm *Moniezia ovisae*. International multidisciplinary Research Journal. 1(12): 01-03.
15. Humbe, et. al. (2012). A report of new mammalian tapeworm *Moniezia osmanabadensis* from *Capra hircus* (L.) at Osmanabad. District (M.S.), India. Journal of experimental science. 3(5): 08-10
16. Kalse A.T. and G.B. Shinde, (1999). On *Moniezia* (*Blanchariezia*) *murhari*, n. sp. (Cestoda :Anoplocephalidae) *Fuhrmann*, 1907) from *Capra hircus* (L.) in (M.S.) India. Rivista Di Parasitologia, Vol XVI (LX) N.1 APRILE 1999
17. Monnig H.O. (1926). Three new helminthes Trans. Ray.Soc.South Africa.13:291-298.
18. Nanware, S. S. (2010). Report on occurrence of *Moniezia* (*Blanchariezia*) *Caprae* Sp. Nov. (Cestoda: Anoplocephalidae) from *Capra hircus* (L.). The Biosphere. 2(1): 27-30.
19. Nanware, S. S., (1999). A new record of *Moniezia* (*Blanchariezia*) *kalavati* n. sp. from *Capra hircus* L. 13 th Nat. Cong. Parasitol. Eb.24-26. 1999. Sou. Abstract no.164, pp. 118.
20. Padwal Nitin and M. N. Kadam, (2011). Report of a new mammalian tapeworm *Moniezia govindae*. Rec Res Sci Tech 3 (2011) 30-33.
21. Patil, S. R. and Shinde G. B., (1997). A new species of the cestode *Moniezia*. (B) *waranaganarensis*, n. sp from Sheep. Riv. Di. Parasit. XIV(LVIII) N-2A: 905-997.
22. Pawar, S. B., (2004). A new cestode *Moniezia* (*Blanchariezia*) *shindei* n. sp. from *Capra hircus* M.S. India. Rivista Di Parasit. XII (LXV) – N 2: 87 – 90.
23. Pokale, S. N., (2004). On a new species of *Moniezia caprai* Blanchard, 1891 (Cestoda: Anoplocephalidae) from *Capra hircus*. Utter Pradesh J. Zool. 24 (3): 285-288.
24. Rawankar, Amol S., Gajanan A. Wagh, and Jayant S. Wadatar (2015). "DNA barcoding and phylogenetic analysis of *Tyto alba*, *Otus bakkamoena* and *Athene brama* from Indian subcontinent." International J. Zoology and Res. (IJZR) 5: 7-14
25. Shaikh Kalim, (2015). Biosystematic studies on *Moniezia* (B) *marathwadensis* sp. nov. parasitic in *Capra hircus* from Aurangabad District, MS India., International Journal of Current Microbiology and Applied Sciences 4(10): 409-416
26. Shinde G.B. Jadhav B.V. and S.S. Kadam, (1985). Two new species of the genus *Moniezia* Blanchard 1891 Rivista Di Parasitologia, Vol.II (XLVI) APRILE 1985
27. Skrjabin, K. J. and R. I. Schulz, (1937). Helminthology Miskow, 2nd Ed. PP. 418.
28. Solunke Ravi, (2015) On a new species of *Moniezia* (Blanchard 1891) (Cestoda: Anoplocephalidae) in *Capra hircus* (L.) from Latur District (M.S.) India. pp. 514-516.
29. Sunita Borde, Amol Thosar, Sapna Lohat, Rahul Khawal and Sushil Jawale, (2017). Diversity of Cestode parasites in Vertebrates from Marathwada region (M.S.) India. International Journal of the Social Research Foundation. IIC-DESW 2017-Special Issue: Vol. No. II. 276-283.
30. Sunita Borde, Ashwini Bhosale and Amol Thosar, (2021). Molecular and Morphological identification of two Cestode Species Parasitizing Sheep and Goats in Aurangabad District (M.S.) India, Natural Volatiles & Essential Oils, 8(4):3225-3340
31. Sunita Borde, Ashwini Bhosale and Amol Thosar, (2022). Molecular and Morphological identification of Cestode parasites from goat in Aurangabad

- district (M.S.) India., Journal of Positive School Psychology (JPSP),8(4):
32. Tat, M. B. and B. V. Jadhav. (2004). A new tapeworm from the host, *Capra hircus* at Beed (Maharashtra) India. Nat. J. Life. Sci. PP. 255-258.
  33. Vishwakarma, sunilkumar, A. M. R. I. T. A. Nigam, and A. T. U. L. Singh. (2016) "Molecular phylogenetic analysis of *Fusarium* isolates causing pokkahboeng disease in sugarcane based on RAPD marker." International Journal of Agricultural Science and Research 6.3: 177-186.
  34. Yamaguti, S. (1959). *Systema Helminthum*, Vol. II, The Cestodes of vertebrates, Interscience Pub. INC, New York London, 1-860.