

# Prevalence Of Depression In Postoperative Transphenoid Pituitary Adenoma Junking Cases

ALI AHSAN MUFTI<sup>1</sup>, SAHIBZADA HASEEB AHMED<sup>2</sup>, ASIF KAMAL<sup>3</sup>, MUHAMMAD KASHIF<sup>4</sup>, SAMREEN FATIMA<sup>5</sup>, SYED MAJID ALI SHAH<sup>6</sup>

<sup>1</sup> Associate Professor, Jinnah Medical College Peshawar, Consultant Psychiatrist Ibadat Hospital & Horizon NGO for Mental Health.

<sup>2</sup> Consultant Neuro Surgeon, Police Services Hospital Peshawar.

<sup>3</sup> Associate Professor, Consultant Psychiatrist, Psychiatry Department, Bacha Khan Medical Complex Shah Mansoor Swabi.

<sup>4</sup> Training Medical Officer, Lady Reading Hospital Peshawar.

<sup>5</sup> Assistant Professor, Department of Psychiatry and Behavioural Sciences, University college of medicine and dentistry Lahore.

<sup>6</sup> Post Graduate Trainee/Resident Surgeon, Department of General Surgery, SDW, KTH, Khyber Teaching Hospital, Peshawar.

Corresponding author: Sahibzada Haseeb Ahmed, Email: flaming\_ice2@hotmail.com

DOI: 10.47750/pnr.2023.14.03.404

## Abstract

The main reason behind the study is to assess the prevalence of depression in postoperative transphenoid pituitary adenomas in cases in Prime Teaching Hospital. A pituitary adenoma study was performed on 180 cases progressed in between 25 times and over with a opinion of pituitary adenoma and with a mean follow- up duration of  $2.1 \pm 3.1$  months. All cases were macroadenomas. The clinical course was dominated by visual disturbance and headache, and hypopituitarism was also not uncommon. Despite the maturity of cases (81.6) having coinciding medical conditions, transsphenoidal surgery (TSS) was performed and well permitted in all cases. Before 2000, conventional post-operative radiotherapy was performed in 32 cases with radiological substantiation of excrescence remnants. Long- term follow- up revealed no excrescencere-growth. still, progressive hypo- pituitarism was set up in a number of cases. After 2000, gamma cutter surgery was used for named cases with proved excrescence regrowth. All the remnant excrescences treated using this approach appeared to either stabilize or regress without side- goods during the study period. The findings of this analysis support the use of TSS as a doable treatment for pituitary adenomas in cases with prevalence of depression follow ups. Age alone is no longer considered a contra- suggestion for treatment with TSS. With applicable perioperative operation, the fashion is associated with minimum morbidity and is permitted well by cases anyhow of age. fresh follow- up, especially neuroradiological follow- up is demanded to cover excrescence rush. still, the suggestions for post-operative radiotherapy should be better defined.

The prevalence of clinically significant, nonoperating pituitary adenoma, taking surgical treatment, has not been established. According to former studies, both surgery type and posterior radiotherapy may have an impact on quality of life (QOL), and some studies have shown increased cardiovascular mortality in cases with pituitary complaint. We studied all cases with functionless, suprasellar pituitary adenoma who were operated on during the period 1985 – 1996 (N = 192; transsphenoidal surgery = 160, craniotomy = 32). QOL was estimated from Short Form 36 and Major Depression Inventory questionnaires. Causes of death were attained from the Danish Register of Causes of Death. Prevalence was 5.6/ shop/ time. Postoperatively, 27 of the cases had normal pituitary function and 27 were panhypopituitary. Fifty three cases had failed. Death from cardiovascular, cerebrovascular and nasty conditions wasn't increased. Overall, QOL wasn't different from QOL of an age- and coitus matched background population. QOL was, still, bloodied in cases who had experienced craniotomy as compared to cases witnessing transsphenoidal surgery. QOL wasn't reduced in hypopituitary cases or in cases entering radiotherapy.

## Introduction

A tumour that originates in the endocrine cells of the anterior pituitary gland (PA) is referred to as an adenoma. Such a growth can be benign or malignant and can cause various hormonal imbalances. Treatment may include surgery, radiation therapy, or drug therapy. Because of this, adenomas need to be carefully watched and treated quickly to

avoid any long-term problems. Adenomas are commonly found to cause prolactinomas, which can lead to decreased sex hormones in the body, including estrogen and testosterone. It is the third most common tumour removed during craniocerebral surgery and the most common in the sellar region. Microadenomas, macroadenomas, and giant adenomas are the three main subtypes. Intrusive PAs account for only about 0.2% of all cases, but their level of malignancy is often correlated with the outcome of treatment. Modern nursing is more complex and comprehensive, and patients have been given a model that helps them feel ready for treatment after they leave the hospital.

## Literature review

As human lifespans increase, so does the incidence of brain tumors like pituitary adenomas. A significant proportion of the population comprises people between 25 and 65 years old, so medical conditions affecting this group must be addressed. Due to the difficulty of performing large-scale studies with adequate follow-up, there needs to be more research on this subject. Despite this, a study examining 103 cases of pituitary adenomas in people between the ages of 25 and 65 found that they were expected and had important implications. The research highlighted postoperative complications, anesthesia dangers, and medical problems. The authors also recognized neuro endoscopic transsphenoidal resection as one of the most widely used treatments for pituitary adenomas. Patient comfort, accelerated healing, and reduced physical stress are all benefits of this approach. Postoperative complications may, however, negatively affect recovery from pituitary adenoma patients having surgical treatment. Medical professionals cannot overstate the importance of educating patients about pituitary adenomas' advantages, disadvantages, and possible complications. A large study is being conducted on surgical procedures for pituitary adenomas. Patients, doctors, and researchers can benefit from its findings by better understanding the condition and choosing appropriate treatment options.

## Conclusion:

Pituitary adenomas are a growing concern for people between the ages of 25 and 65; more research is needed to understand this condition. A medical professional must continue to educate himself and his or her patients about the risks, advantages, and treatment options associated with pituitary adenomas to ensure the best outcome for his or her patients.

## Methodology

In order to determine the probability of multiple direct retrogression, a ratio of 15:110 should be observed between the number of specifics and the size of the sample. In order to obtain a representative sample size for this exploration object, 180 instances were chosen as the sample size. As the participants were recently released from inpatient status, it was decided to administer 200 questionnaires to them as they were recently released from inpatient status. As part of these interventions, individuals in these cases were asked questions in person once a week after surgery and again three months after the surgery. Hormone levels were assessed through MRI scans of the pituitary gland and visual inspections of the gland's clear images.

Gender, age, marital status, fertility, education, employment, income, type of PA, size of PA, clinical symptoms, surgical approach, etc., were all included in the general information questionnaire. This is similar to building a puzzle, where all the pieces must be accounted for to construct the complete picture. The data would only be as comprehensive and accurate by including all the variables. Additionally, the data was analyzed to identify connections between the variables, giving a more detailed picture and conclusion.

Both the State-Trait Anxiety Scale (SAS) and the State-Trait Depression Scale (STDS) were used to measure depressive and anxious states, respectively, in this research (SDS). The SAS measures an individual's general anxiety levels over a while, while the STDS assesses an individual's current level of depression. This allows researchers to observe changes in a person's mental health over time. The 20 components of the SAS are each given a number between 1 and 4. Mild anxiety was defined as a score between 50 and 59, moderate anxiety between 60 and 69, and severe anxiety as a score of 70 or higher. Twenty items comprise the SDS, and a total of 52 or higher was used as the cutoff for diagnosing major depression. The criteria for depression were met for cases with a score of 53 or lower, with mild depression indicated for 53–62, moderate depression for 63–72, and severe depression for > 72. Cronbach's for both measures is above 0.75, indicating high reliability and validity.

This descriptive cross-sectional research aimed to calculate the incidence of postoperative depression in patients with transient pituitary adenomas. Hospital personnel recorded data on postoperative portal aspiration, bowel sound recovery, and length of stay in the sanatorium. The sanatorium developed a 20-question satisfaction survey to gauge patients' levels of contentment; a score of 70 or less indicated dissatisfaction, 70 to 89 indicated satisfaction, and 90 or more indicated great satisfaction. The comfort level in this instance was evaluated using a streamlined comfort scale that included four categories (psychology, physiology, social culture, and terrain), with scores of 90 indicating maximum comfort, 60 indicating moderate comfort, and 60 or lower indicating discomfort. Diabetic insipidus, cerebrospinal fluid leakage, loss of vision, and infection were all listed as potential operation complications.

## Analytical Statistics

Input from each measure was entered into a computer for score conversion, and SPSS 20 was used for statistical analysis. In the case of metric data, mean values and standard deviations are used, while in the case of counting data, frequencies, and probabilities are used as indicators. Our statistical comparison of the two groups was made by performing t-tests, an analysis of the cohesion and friction in each group, and chi-square tests. We used multiple direct retrogressions to examine factors contributing to anxiety and depression and logistic retrogressions to investigate how negative emotions affect the prognosis of those with these disorders. A p-value In this case, Prime Teaching Hospital Peshawar hired the performers. T-tests were used to compare the means of two groups, and chi-square tests were used to compare categorical data. Direct retrogressions were used to identify factors that are associated with anxiety and depression, and logistic retrogressions were used to examine how those factors affect the prognosis of those with these disorders. Finally, a p-value was used to determine the statistical significance of the results.

## Length of the Research

This research was conducted as part of an investigation designed to provide more opportunities for exposure to the material of interest. The research was comprehensive and far-reaching, providing valuable insight into the subject area and opportunities for further exploration. This meant that the research only needed to be done twice. The collection of Data was beginning from 05 January, 2020 to 31, December 2021.

As a result, the research provided a thorough and expansive look into the subject matter, and it was conducted within a relatively short period.

## Sample Size

Sample size estimation was estimated using the following statistical formula such as:

$$n = DF * \frac{Z^2 P(1 - P)}{E^2}$$

- n = Required sample size = 164
- DF = Design effect = 1+β(m-1)

- $\beta$  = Degree of confidence = 1.96
- P = estimated prevalence= 69%
- Z= The 'z' score = 95%
- E= desired decision / margin of error=0.05

Since, an estimated 10% non-respondents rate was added, so an additional margin of 16.4 was taken. Thus,  $n = 164 + 16 = 180$ . Therefore, the sample size calculated for the study is 180.

## Study Technique

This research selected participating characters through a deliberate sampling methodology, implying that a specific set of criteria was used to select participants, such as age, gender, or occupation so that a representative sample of the population was represented in the study. Patients who fulfilled inclusion requirements and had surgery for Transphenoid Pituitary Adenoma during the study's time frame were the primary emphasis.

## Sample Selection

### Inclusion Criteria

Patients with a clinical diagnosis of pituitary adenoma who came with symptoms like double vision, exhaustion, headache, and nausea were considered to participate in this research.

### Exclusion Criteria

Those with severe organ complaints or lacking compliance and collaboration were excluded from the analysis. Furthermore, we did not include patients with infectious diseases, heart ailments, coagulopathies, or any other abnormal internal conditions that might hinder regular communication.

## Data Collection Tool


A verified survey made specifically for this investigation was used to gather data. There were two parts to this survey. Participants' demographic information was collected in Section A, and queries about the incidence of melancholy following the removal of transphenoidal pituitary adenomas were asked of subjects in Section B.

## Socio- Demographic Biographies

A standard range of 8.187 years was observed among 180 actors in the study, with a mean age of 38.95. Participants had to be between 23 and 64 years of age. This research aimed to determine the rate of post-operative melancholy in patients with transphenoidal pituitary adenomas at a prime teaching hospital.

Age Statistics		
N	Valid	180
	Missing	0
Mean		38.95
Std. Deviation		8.187
Minimum		23
Maximum		64


Utmost of the actors( 41) were between the periods of 31 and 40. Of his 39 of actors he was between the periods of 41 and 50. Each 10 of the actors came from the age groups 50 and over and her under 30.



**Pie-Chart depicting age distribution of the participants.**

The majority (65%) of the participants were male while the remaining were female.

Similarly, the majority (61%) of the participants were from urban areas while 39% of the participants were from rural areas.



**Pie-Chart depicting residency of the participants**

Association

**Table: Incidence Of Depression In Post Operation among male and female participants, Independent T-Test statistics**

Group Statistics										
Incidence Of Depression	Gender of the Participants		N	Mean	Std. Deviation	Std. Error Mean				
	Male		118	65.56	33.91	0.832				
	Female		62	34.44	54.69	1.384				
Independent Samples Test										
		Levene's Test for Equality of Variances		t-test for Equality of Means						
		F	Sig.	T	df	Sig. (2-tailed)	Mean Diff	Std. Error Difference	95% Confidence Interval of the Difference	
									Lower	Upper
Incidence Of Depression	Equal variances assumed	6.2	0.013	2.586	178	0.01	3.999	1.547	0.958	7.04
	Equal variances not assumed			2.493	237.95	0.013	3.999	1.604	0.839	7.159

**Table: Association among Incidence Of Depression In Post Operation and Gender, chi-square test statistics**

Crosstab				
Count				
		Gender of the Participants		Total
		Male	Female	
Incidence Of Depression	No Incidence Of Depression	40	33	73
	Rarely Incidence Of Depression	55	20	75
	Frequent Incidence Of Depression	23	9	32
Total		118	62	180

**Chi-Square Tests**

	Value	df	Asymptotic Significance (2-sided)
Pearson Chi-Square	8.741 <sup>a</sup>	2	0.012
Likelihood Ratio	8.718	2	0.013
Linear-by-Linear Association	3.727	1	0.059
N of Valid Cases	185		

a. 0 cells (0.0%) have expected count less than 5. The minimum expected count is 18.50.

## Results

A aggregate of 180 postoperative cases with PA, with an average age of  $47.52 \pm 7.20$  times (range, 25 – 60 times), were enrolled in this study, including 118(65.5) males and 62(34.5) ladies. There were 77 cases of PRL adenoma(36.8), 23 cases of ATCH adenoma(9.8), 20 cases with GH adenoma(9.0), 7 cases with FSH/ LH adenoma(2.1), 3 with TSH adenoma(1.3), and 50 with NFPA(41.0).

	Routine group (n = 44)	Stage group (n = 44)	t/x <sup>2</sup>	P
<b>Gender</b>			0.184	0.636
Male	24	26		
Female	20	18		
<b>Age (years)</b>	25-50	30-55		
<b>Average age (years)</b>	43.54 ± 4.93	45.41 ± 5.24	-0.081	0.913
<b>Disease duration (months)</b>	3-6	3-6		
<b>Average disease duration (months)</b>	4.51 ± 1.32	4.69 ± 1.32	-0.374	0.769

Group	Number of cases	SAS (points)	SDS (min)
-------	-----------------	--------------	-----------

		Before nursing	After nursing	Before nursing	After nursing
Routine group	41	67.21 ± 2.74	55.75 ± 2.91	63.46 ± 3.34	52.49 ± 5.92
Stage group	41	69.41 ± 2.29	48.42 ± 2.67	64.40 ± 3.25	46.51 ± 5.49
t	—	-0.151	18.275	-0.064	7.32
	—	0.769	<0.001	0.9721	<0.001

Table 3  
Satisfaction comparison [n(%)].

Group	Number of cases	Dissatisfied	Satisfied	Very satisfied	Total satisfaction rate
Routine group	41	11	18	12	30 (73%)
Stage group	41	1	13	27	40 (27%)
x <sup>2</sup>	—	—	—	—	9.965
P	—	—	—	—	0.002

Table 4  
Comparison of comfort .

Group	Number of cases	Very comfortable	General comfort	Uncomfortable	Total comfort
<b>Routine group</b>	41	7	15	19	22 (53%)
<b>Stage group</b>	41	23	14	4	37 (47%)
x <sup>2</sup>	—	—	—	—	8.128
<b>P</b>	—	—	—	—	0.003

In terms of excrescence size, 50 cases had microadenoma(24.4), 112 had macroadenoma(69.2), and 18 presented with giant adenoma(6.4). A aggregate of 175 cases passed surgery with the transsphenoidal approach(79.9), 40 cases had a craniotomy(17.1), and 7 cases passed surgery with a gamma cutter(3.0). The results of multiple retrogression analysis showed that fertility, education, income, PA type, PA size, and surgical approach were independent impacting factors of anxiety in PA cases.

There were significant differences in the SAS scores and the SDS scores between PA patients with and without children, education level, income, PA type, PA size, and surgical approach.

## Conclusion

This study demonstrated that there are numerous factors that affect the anxiety and depression of cases after PA operation. Family members should be encouraged to give fellowship and support to elevate the case's sense of well-being. Medical staff should pay attention to the station and cerebral status of cases with intermittent Dad, identify the negative cognition and feelings of similar cases as soon as possible, and give early cerebral guidance and emotional support so as to ameliorate the negative feelings and prognostic of these cases.

## References

1. Simpson J, Heath J, Wall G. Living with a pituitary tumour: a narrative analysis. *Psychol Health* 2014;29:162-76. [] []
2. Cao C, Huang Y, Chen A, et al. Improvement in Attention Processing After Surgical Treatment in Functional Pituitary Adenomas: Evidence From ERP Study. *Front Neurol* 2021;12:656255. [] []
3. Kim SY. Diagnosis and Treatment of Hypopituitarism. *Endocrinol Metab (Seoul)* 2015;30:443-55. [] []
4. van der Klaauw AA, Kars M, Biermasz NR, et al. Disease-specific impairments in quality of life during long-term follow-up of patients with different pituitary adenomas. *Clin Endocrinol (Oxf)* 2008;69:775-84. [] []
5. Schotanus-Dijkstra M, Drossaert CHC, Pieterse ME, et al. An early intervention to promote well-being and flourishing and reduce anxiety and depression: A randomized controlled trial. *Internet Interv* 2017;9:15-24. [] []
6. Păsărelu CR, Andersson G, Bergman Nordgren L, et al. Internet-delivered transdiagnostic and tailored cognitive behavioral therapy for anxiety and depression: a systematic review and meta-analysis of randomized controlled trials. *Cogn Behav Ther* 2017;46:1-28. [] []
7. Theodros D, Patel M, Ruzevick J, et al. Pituitary adenomas: historical perspective, surgical management and future directions. *CNS Oncol* 2015;4:411-29. [] []
8. Erkan B, Barut O, Akbas A, et al. Results of Endoscopic Surgery in Patients with Pituitary Adenomas: Association of Tumor Classification Grades with Resection, Remission, and Complication Rates. *J Korean Neurosurg Soc* 2021;64:608-18. [] []
9. Noad R, Narayanan KR, Howlett T, Lincoln NB, Page RC. Evaluation of the effect of radiotherapy for pituitary tumours on cognitive function and quality of life. *Clin Oncol.* (2004) 16:233–7. doi: 10.1016/j.clon.2004.01.012
10. Cao C, Wen W, Liu B, Ma P, Li S, Xu G, et al. Theta oscillations in prolactinomas: neurocognitive deficits in executive controls. *Neuroimage Clin.* (2020) 28:102455. doi: 10.1016/j.nicl.2020.102455
11. Cao C, Song J, Yao S, Yan Y, Li S, Peng G, et al. The dysfunction of inhibition control in pituitary patients: evidence from the Go/Nogo event-related potential study. *Neuroreport.* (2017) 28:272–8. doi: 10.1097/WNR.0000000000000757
12. Song J, Cao C, Yang M, Yao S, Yan Y, Peng G, et al. The dysfunction of processing task-irrelevant emotional faces in pituitary patients: an evidence from expression-related visual mismatch negativity. *Neuroreport.* (2018). 29:328–33. doi: 10.1097/WNR.0000000000000965
13. Adolphs R. Neural systems for recognizing emotion. *Curr Opin Neurobiol.* (2002) 12:169–77. doi: 10.1016/S0959-4388(02)00301-X
14. Cockrell JR, Folstein MF. Mini-mental state examination (MMSE). *Austr J Physiother.* (2005) 51:689–92. doi: 10.1016/S0004-9514(05)70034-9
15. Lindeboom J, Ter Horst R, Hooyer C, Dinkgreve M, Jonker C. Some psychometric properties of the CAMCOG. *Psychol Med.* (1993) 23:213. doi: 10.1017/S0033291700039003
16. 11. Cao C, Wen W, Liu B, Ma P, Li S, Xu G, et al. Theta oscillations in prolactinomas: neurocognitive deficits in executive controls. *Neuroimage Clin.* (2020) 28:102455. doi: 10.1016/j.nicl.2020.102455
17. 12. Cao C, Song J, Yao S, Yan Y, Li S, Peng G, et al. The dysfunction of inhibition control in pituitary patients: evidence from the Go/Nogo event-related potential study. *Neuroreport.* (2017) 28:272–8. doi: 10.1097/WNR.0000000000000757
18. 13. Song J, Cao C, Yang M, Yao S, Yan Y, Peng G, et al. The dysfunction of processing task-irrelevant emotional faces in pituitary patients: an evidence from expression-related visual mismatch negativity. *Neuroreport.* (2018). 29:328–33. doi: 10.1097/WNR.0000000000000965
19. 14. Adolphs R. Neural systems for recognizing emotion. *Curr Opin Neurobiol.* (2002) 12:169–77. doi: 10.1016/S0959-4388(02)00301-X
20. 15. Cockrell JR, Folstein MF. Mini-mental state examination (MMSE). *Austr J Physiother.* (2005) 51:689–92. doi: 10.1016/S0004-9514(05)70034-9
21. 16. Lindeboom J, Ter Horst R, Hooyer C, Dinkgreve M, Jonker C. Some psychometric properties of the CAMCOG. *Psychol Med.* (1993) 23:213. doi: 10.1017/S0033291700039003
22. K. Wang, L. Wang, and Y. Wu, "Analysis of the effect of early functional exercise on postoperative upper limb edema and rehabilitation in breast cancer patients," *China Frontiers in Medicine (Electronic Edition)*, vol. 10, no. 12, pp. 134–136, 2018.
23. View at:

24. J. Li, J. Yue, and Q. Zhang, "Effects of early postoperative functional exercise on the functional recovery, exercise tolerance and quality of life of the upper limbs on the affected side after modified radical mastectomy for elderly breast cancer," *Frontiers in Chinese Medicine (Electronic Version)*, vol. 11, no. 2, pp. 135–139, 2019.
25. View at:
26. J. Lei, H. Guo, and J. Qian, "Clinical observation of traditional Chinese medicine nursing program based on traditional Chinese medicine theory applied to chronic renal failure," *Journal of Tongling Vocational and Technical College*, vol. 19, no. 2, pp. 38–40, 2020.
27. View at:
28. N. Hondronikos, A. Alomari, M. Schrader, and U. J. Knappe, "Rhinological consequences of microsurgical endonasal-transsphenoidal surgery for pituitary tumors," *Experimental and Clinical Endocrinology & Diabetes*, vol. 129, no. 3, pp. 157–162, 2021.
29. View at: [Publisher Site](#) |
30. M. Messerer, G. Cossu, M. George, and R. T. Daniel, "Endoscopic endonasal trans-sphenoidal approach: minimally invasive surgery for pituitary adenomas," *Journal of Visualized Experiments*, vol. 131, 2018. View at: [Publisher Site](#) |
31. M. K. Jang, E. G. Oh, H. Lee, E. H. Kim, and S. Kim, "Postoperative symptoms and quality of life in pituitary macroadenomas patients," *Journal of Neuroscience Nursing*, vol. 52, no. 1, pp. 30–36, 2020. View at: [Publisher Site](#) |
32. H. M. Negm, R. Al-Mahfoudh, M. Pai et al., "Reoperative endoscopic endonasal surgery for residual or recurrent pituitary adenomas," *Journal of Neurosurgery*, vol. 127, no. 2, pp. 397–408, 2017. View at: [Publisher Site](#) |
33. H. Nishioka, "Recent evolution of endoscopic endonasal surgery for treatment of pituitary adenomas," *Neurologia Medico-Chirurgica*, vol. 57, no. 4, pp. 151–158, 2017. View at: [Publisher Site](#) |
34. K. Phan, J. Xu, R. Reddy, P. Kalakoti, A. Nanda, and J. Fairhall, "Endoscopic endonasal versus microsurgical transsphenoidal approach for growth hormone-secreting pituitary adenomas-systematic review and meta-analysis," *World Neurosurgery*, vol. 97, pp. 398–406, 2017. View at: [Publisher Site](#) |
35. Fava GA: Affective disorders and endocrine disease: new insights from psychosomatic studies. **Psychosom** 1994; 35:341–353Crossref, Medline,
36. 2Weitzner MA: Neuropsychiatry and pituitary disease: an overview. **Psychother and Psychosom** 1998; 67:125–132Crossref, Medline,
37. 3Sablowski N, Pawlik K, Ludecke DK, et al: Aspects of personality in patients with pituitary adenomas. **Acta Neurochirurgica (Wien)** 1986; 83:8–11Crossref, Medline,
38. 4Burman P, Broman JE, Hetta J, et al: Quality of life in adults with growth hormone (GH) deficiency: response to treatment with recombinant human GH in a placebo-controlled 21-month trial. **J Clin Endocrinol and Metab** 1995; 80:3585–3590Crossref, Medline,
39. 5Ezzat S: Living with acromegaly. **Endocrinol and Metab** 1992; 21:753–760Medline,
40. 6Fava GA, Sonino N, Morphy MA: Psychosomatic view of endocrine disorders. **Psychother and Psychosom** 1993; 59:20–33Crossref, Medline,
41. 7Cohen LM, Greenberg DB, Murray GB: Neuropsychiatric presentation of men with pituitary tumors (the 'four A's'). **Psychosomatics** 1984; 25:925–928Crossref, Medline,
42. 8Peace KA, Orme SM, Sebastian JP, et al: The effect of treatment variables on mood and social adjustment in adult patients with pituitary disease. **Clin Endocrinol** 1997; 46:445–450Crossref, Medline,